

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: **90107640.6**

(51) Int. Cl.⁵: **A61K 31/415**

(22) Date of filing: **10.03.86**

This application was filed on 23 - 04 - 1990 as a divisional application to the application mentioned under INID code 60.

(30) Priority: **14.03.85 GB 8506642**
09.04.85 GB 8509039
23.12.85 GB 8531614

(43) Date of publication of application:
05.09.90 Bulletin 90/36

(60) Publication number of the earlier application in accordance with Art.76 EPC: **0 201 165**

(84) Designated Contracting States:
BE CH DE FR GB IT LI NL


(71) Applicant: **Beecham Group p.l.c.**
SB House Great West Road
Brentford Middlesex TW8 9BD(GB)

(72) Inventor: **Wootton, Gordon, Beecham**
Pharmaceuticals
Coldharbour Road, The Pinnacles
Harlow, Essex CM19 5AD(GB)
 Inventor: **Sanger, Gareth John, Beecham**
Pharmaceuticals
Coldharbour Road, The Pinnacles
Harlow, Essex CM19 5AD(GB)

(74) Representative: **Jones, Pauline et al**
SmithKline Beecham, Corporate Patents,
Great Burgh, Yew Tree Bottom Road
Epsom, Surrey KT18 5XQ(GB)

(54) **Medicaments for the treatment of emesis.**

(57) The use of a compound of formula (II) or a pharmaceutically acceptable salt thereof:



wherein

R₁₀ is hydrogen, C₁₋₁₀ alkyl, C₃₋₇ cycloalkyl, C₃₋₆ alkenyl, phenyl or phenyl-C₁₋₃ alkyl; and one of the groups represented by R₁₁, R₁₂ and R₁₃ is hydrogen, C₁₋₆ alkyl, C₃₋₇ cycloalkyl, C₂₋₆ alkenyl or phenyl-C₁₋₃ alkyl and each of the other groups, which may be the same or different, is hydrogen or C₁₋₆ alkyl; in the manufacture of a medicament for use in the treatment of cytotoxic agent or radiation induced nausea and vomiting, or motion sickness.

MEDICAMENTS FOR THE TREATMENT OF EMESIS

The present invention relates to a method of treatment of emesis, such as cytotoxic agent or radiation induced nausea and vomiting or motion sickness; anxiety; and irritable bowel syndrome (IBS); in mammals including humans, and to the use of compounds in the preparation of a medicament for the treatment of such conditions.


5 U.K. Patent Applications, 2100259, 2125398 and 2152049 describe classes of compounds which are aryl amides and esters having an azabicyclic side chain which are 5-HT receptor antagonists useful in the treatment of migraine. U.K. Patent Application 2153821 describes a class of tetrahydrocarbazolones which are also 5-HT receptor antagonists.

10 It has now been discovered that these classes of compounds and related compounds block the emetic response induced by cytotoxic agents such as cisplatin and also by radiation treatment. The compounds are therefore of use in the treatment of nausea and vomiting associated with cancer therapy. The compounds are also of potential use in the treatment of motion sickness, anxiety and irritable bowel syndrome.

15 Accordingly, the present invention provides a method of treatment of emesis, anxiety and/or IBS in mammals, including humans, which method comprises administering an effective amount of a compound of formula (I) or (II) or a pharmaceutically acceptable salt thereof;

Ar-CO-Y-Z (I)


wherein Ar is a group of formula (a):



(a)

wherein:

30 R₁ and R₂ are independently selected from hydrogen, halogen, C₁₋₆ alkyl, C₁₋₄ alkoxy, hydroxy, amino optionally substituted by one or two C₁₋₄ alkyl groups, thiol, C₁₋₄ alkylthio; X is CH₂, NR₃, -O- or -S- wherein R₃ is hydrogen, C₁₋₄ alkyl, C₃₋₅ alkenyl, phenyl or phenyl C₁₋₄ alkyl; or Ar is a group of formula (b):




(b)

45 wherein R₄ to R₇ are independently hydrogen, halogen, C₁₋₆ alkyl, C₁₋₆ alkoxy, hydroxy, amino optionally substituted by one or two C₁₋₄ alkyl, by C₁₋₄ alkanoylamino or pyrrolyl, one of R₄ to R₇ being other than hydrogen;

Y is -O- or -NH-; and

Z is a group of formula (c), (d) or (e):

50




(c)


wherein

n is 2, 3 or 4;

R₈ is hydrogen, C₁₋₇ alkyl, C₃₋₅ alkenyl or phenyl C₁₋₄ alkyl optionally substituted by one or two halogen, C₁₋₄ alkoxy or C₁₋₄ alkyl;



(d)




(e)

wherein:

p is 1, 2 or 3; and

R₉ is as defined above for R₈;

and with the proviso that, when Ar is of formula (b) and Y is -NH-, Z is a group of formula (d) or (e);



(II)

wherein

R₁₀ is hydrogen, C₁₋₁₀ alkyl, C₃₋₇ cycloalkyl, C₃₋₆ alkenyl, phenyl or phenyl-C₁₋₃ alkyl; and one of the groups represented by R₁₁, R₁₂ and R₁₃ is hydrogen, C₁₋₆ alkyl, C₃₋₇ cycloalkyl, C₂₋₆ alkenyl or phenyl-C₁₋₃ alkyl and each of the other groups, which may be the same or different, is hydrogen or C₁₋₆ alkyl.

In formula (I):

When Ar is a group of formula (a), suitable values for R₁ include hydrogen, chloro, bromo, methoxy, ethoxy, n- and iso-propoxy, amino optionally substituted by one or two methyl groups, thiol, methylthio, ethylthio, n- and iso-propylthio. Preferably R₁ is hydrogen.

In the group of formula (a) the carbonyl side chain may be attached to the ring carbon atom in positions 2, 3, 4, 5, 6 or 7 of the nucleus, but preferably in position 3, 4 and 5. Most preferably the carbonyl group is

attached to the ring containing X especially in position 3. Preferably X is then NR_3 as defined.

R_1 is attached to the ring carbon atom in position 4, 5, 6 or 7 of the nucleus, preferably position 5 and R_2 is attached to the ring carbon atom in position 2 or 3 of the nucleus. Tautomers are also covered by formula 1 e.g. when R_2 is hydroxy or thiol in the 2 position.

5 R_3 is usually hydrogen or alkyl.

When Ar is a group of formula (b) there is a first group of compounds wherein:

R_4 is halogen, C_{1-4} alkylamino or C_{1-4} alkoxy;

R_5 is hydrogen or halogen;

R_6 is hydrogen, amino, nitro, (C_{1-4}) alkylamino, or $\text{di}(\text{C}_{1-4})$ alkylamino, halogen or 1-pyrrolyl;

10 R_7 is hydrogen or halogen, and

preferably R_8 is other than hydrogen, halogen or pyrrolyl.

When Ar is a group of formula (b) there is a second preferred group of compounds wherein:

Examples of R_5 include halo such as chloro, C_{1-6} alkyl such as methyl, and C_{1-6} alkoxy, such as methoxy, preferably chloro or methyl; examples of R_6 include hydrogen, halo, such as chloro, hydroxy, C_{1-6} alkyl
15 such as methyl and C_{1-6} alkoxy, such as methoxy, preferably hydrogen or chloro; examples of R_7 include hydrogen, halo such as chloro, C_{1-6} alkyl such as methyl and C_{1-6} alkoxy, such as methoxy, preferably hydrogen, methyl or chloro; R_4 is then often hydrogen.

Examples of R_8 in formula (c) and R_9 in formula (e) when C_{1-7} alkyl include as groups of interest C_{1-3} alkyl such as methyl, ethyl and *n*- and *iso*-propyl. Within C_{1-7} alkyl, C_{4-7} alkyl are also of interest,
20 especially those of the formula $(\text{CH}_2)_u\text{R}_{14}$ wherein *u* is 1 or 2 and R_{14} is a secondary or tertiary C_{3-6} group. Examples of C_{4-7} alkyl include *n*-, *sec*- and *tert*-butyl, *n*-pentyl, *n*-heptyl, and *iso*-butyl, 3-methylbutyl, and *tert*-butylmethyl. R_8 or R_9 is preferably methyl.

Examples of R_8 or R_9 when phenyl C_{1-4} alkyl include benzyl optionally substituted by hydrogen, chloro, bromo, CF_3 , methoxy, ethoxy, *n*- or *iso*-propoxy, methyl, ethyl, *n*- or *iso*-propyl.

25 *n* is preferably 2 or 3, most preferably 2.

p is often 1 or 2, preferably 2.

In formula (II):

It will be understood that when R_{10} represents a C_{3-6} alkenyl group, the double bond may not be adjacent to the nitrogen atom.

30 The alkyl groups represented by R_{10} , R_{11} , R_{12} and R_{13} may be for example, methyl, ethyl, propyl, prop-2-yl, butyl, but-2-yl, methylprop-2-yl, pent-3-yl or hexyl.

An alkenyl group may be, for example, a propenyl group.

A phenyl- C_{1-3} alkyl group may be for example, a benzyl, phenethyl or 3-phenylpropyl group. A cycloalkyl group may be, for example, a cyclopentyl, cyclohexyl or cycloheptyl group.

35 The pharmaceutically acceptable salts of the compounds of the formula (I) and (II) include acid addition salts with conventional acids such as hydrochloric, hydrobromic, boric, phosphoric, sulphuric acids and pharmaceutically acceptable organic acids such as acetic, tartaric, maleic, citric, succinic, benzoic, ascorbic, methanesulphonic, α -keto glutaric, α -glycerophosphoric, and glucose-1-phosphoric acids.

40 The pharmaceutically acceptable salts of the compounds of the formula (I) and (II) are usually acid addition salts with acids such as hydrochloric, hydrobromic, phosphoric, sulphuric, citric, tartaric, lactic and acetic acid.

Preferably the acid addition salt is the hydrochloride salt.

45 Examples of pharmaceutically acceptable salts include quaternary derivatives of the compounds of formula (I) include the compounds quaternised by compounds such as $\text{R}_{15}\text{-T}$ wherein R_{15} is C_{1-6} alkyl, phenyl- C_{1-6} alkyl or C_{5-7} cycloalkyl, and T is a radical corresponding to an anion of an acid. Suitable examples of R_{15} include methyl, ethyl and *n*- and *iso*-propyl; and benzyl and phenethyl. Suitable examples of T include halide such as chloride, bromide and iodide.

Examples of pharmaceutically acceptable salts of the compounds of formula (I) also include internal salts such as pharmaceutically acceptable N-oxides.


50 The compounds of the formula (I) and (II) and, their pharmaceutically acceptable salts may also form pharmaceutically acceptable solvates, such as hydrates, which are included whenever a compound of formula (I) or (II) is herein referred to.

It will of course be realised that some of the compounds of the formula (I) have chiral or prochiral centres, and thus are capable of existing in a number of stereoisomeric forms, including enantiomers. The
55 invention extends to each of these stereoisomeric forms (including enantiomers), and to mixtures thereof (including racemates). The different stereoisomeric forms may be separated one from the other by the usual methods.

In formula (I) when Z is a group of formula c), Y may be in an *exo* or *endo* configuration with respect to


the azabicyclic side chain; the endo configuration is preferred.

A group of compounds within formula (I) is of formula (IA):



15 wherein Z¹ is a group of formula (c) and the remaining variables are as defined in formula (I).


A further group of compounds within formula (I) is of formula (IB):



30 wherein R₅¹ is halo, C₁₋₆ alkyl or C₁₋₆ alkoxy;
R₆¹ is hydrogen or C₁₋₆ alkoxy;

R₇¹ is halo, C₁₋₆ alkyl or C₁₋₆ alkoxy; and Z¹ is as defined in formula (IA).

35 There is another group of compounds within formula (I) of formula (IC):




50 wherein Z² is a group of formula (d) and the remaining variables are as hereinbefore defined.

There is yet another group of compounds within formula (I) of formula (ID):

55

5

10




wherein the variable groups are as hereinbefore defined.

15

There is a further group of compounds within formula (I) of formula (IE):

20

25




wherein Z³ is a group of formula (e) and the remaining variables are as hereinbefore defined.

30

There is a yet further group of compounds within formula (I) of formula (IF):

35

40



45

wherein the variable groups are as hereinbefore defined.

Suitable and preferred values for the variables in formulae (IA) to (IF) are as described for the corresponding variables under formula (I).

A preferred class of compounds represented by the general formula (II) is that wherein R₁₀ represents a hydrogen atom or a C₁₋₆ alkyl, C₃₋₆ cycloalkyl or C₃₋₆ alkenyl group.


50

Another preferred class of compounds represented by the general formula (II) is that wherein one of the groups represented by R₁₀, R₁₁ and R₁₂ represents a C₁₋₃ alkyl, C₃₋₆ cycloalkyl or C₃₋₆ alkenyl group and each of the other two groups, which may be the same or different, represents a hydrogen atom or a C₁₋₃ alkyl group.

55

A further preferred class of compounds represented by the general formula (II) is that wherein R₁₀ represents a hydrogen atom and R₁₂ and R₁₃ represents a C₁₋₃ alkyl group or R₁₀ represents a C₁₋₃ alkyl group and both R₁₂ and R₁₃ represent hydrogen atoms.

A particularly, preferred class of compounds within formula (II) is that represented by the formula (IIA):



(IIA)

wherein R_{10}^1 represents a hydrogen atom or a methyl, ethyl, propyl, prop-2-yl, prop-2-enyl or cyclopentyl group; R_{12}^1 represents a hydrogen atom; and either R_{11}^1 represents a methyl, ethyl, propyl or prop-2-yl group and R_{13}^1 represents a hydrogen atom or R_{11}^1 represents a hydrogen atom and R_{13}^1 represents a methyl or ethyl group. Preferably R_{10}^1 is methyl and R_{12}^1 and R_{13}^1 are both hydrogen.

Compounds of the formula (I) and (II) may be prepared as described in the aforementioned U.K. patent applications, the subject matter of which are incorporated herein by reference, or by analogous methods thereto.

Examples of cytotoxic agents include those routinely used in cancer chemotherapy, such as cisplatin, doxorubicin, cyclophosphamide, particularly cisplatin.

The administration of the compound of formula (I) or (II), or a pharmaceutically acceptable salt thereof may be by way of oral or parenteral administration.

An amount effective to treat the disorders hereinbefore described depends on the relative efficacies of the compounds of the invention, the nature and severity of the disorder being treated and the weight of the mammal. However, a unit dose for a 70kg adult will normally contain 0.5 to 100mg for example 1 to 500mg, of the compound of formula (I) or (II), or a pharmaceutically acceptable salt thereof. Unit doses may be administered once or more than once a day, for example, 2, 3 or 4 times a day, more usually 1 to 3 times a day, that is in the range of approximately 0.001 to 50mg/kg/day, more usually 0.002 to 25mg/kg/day. No adverse toxicological effects are indicated at any of the aforementioned dosage ranges.

It is preferred that the compound of formula (I) or (II) or a pharmaceutically acceptable salt thereof is administered in the form of a unit dose pharmaceutical composition in which is combined with a pharmaceutically acceptable carrier.

Such compositions are prepared by admixture and are suitably adapted for oral or parenteral administration, and as such may be in the form of tablets, capsules, oral liquid preparations, powders, granules, lozenges, reconstitutable powders, injectable and infusable solutions or suspensions or suppositories. Orally administrable compositions are preferred, since they are more convenient for general use.

Tablets and capsules for oral administration are usually presented in a unit dose, and contain conventional excipients such as binding agents, fillers, diluents, tableting agents, lubricants, disintegrants, colourants, flavourings, and wetting agents. The tablets may be coated according to well known methods in the art, for example with an enteric coating.

Suitable fillers for use include cellulose, mannitol, lactose and other similar agents. Suitable disintegrants include starch, polyvinylpyrrolidone and starch derivatives such as sodium starch glycolate. Suitable lubricants include, for example, magnesium stearate.

Suitable pharmaceutically acceptable wetting agents include sodium lauryl sulphate. Oral liquid preparations may be in the form of, for example, aqueous or oily suspensions, solutions, emulsions, syrups, or elixirs, or may be presented as a dry product for reconstitution with water or other suitable vehicle before use. Such liquid preparations may contain conventional additives such as suspending agents, for example sorbitol, syrup, methyl cellulose, gelatin, hydroxyethylcellulose, carboxymethylcellulose, aluminium stearate gel or hydrogenated edible fats, emulsifying agents, for example lecithin, sorbitan monooleate, or acacia; non-aqueous vehicles (which may include edible oils), for example, almond oil, fractionated coconut oil, oily esters such as esters of glycerine, propylene glycol, or ethyl alcohol; preservatives, for example methyl or propyl p-hydroxybenzoate or sorbic acid, and if desired conventional flavouring or colouring agents.

Oral liquid preparations are usually in the form of aqueous or oily suspensions, solutions, emulsions, syrups, or elixirs or are presented as a dry product for reconstitution with water or other suitable vehicle before use. Such liquid preparations may contain conventional additives such as suspending agents, emulsifying agents, non-aqueous vehicles (which may include edible oils), preservatives, and flavouring or colouring agents.

The oral compositions may be prepared by conventional methods of blending, filling or tableting. Repeated blending operations may be used to distribute the active agent throughout those compositions

employing large quantities of fillers. Such operations are, of course, conventional in the art.

For parenteral administration, fluid unit dose forms are prepared containing a compound of the present invention and a sterile vehicle. The compound, depending on the vehicle and the concentration, can be either suspended or dissolved. Parenteral solutions are normally prepared by dissolving the compound in a vehicle and filter sterilising before filling into a suitable vial or ampoule and sealing. Advantageously, adjuvants such as a local anaesthetic, preservatives and buffering agents are also dissolved in the vehicle. To enhance the stability, the composition can be frozen after filling into the vial and the water removed under vacuum.

Parenteral suspensions are prepared in substantially the same manner except that the compound is suspended in the vehicle instead of being dissolved and sterilised by exposure of ethylene oxide before suspending in the sterile vehicle. Advantageously, a surfactant or wetting agent is included in the composition to facilitate uniform distribution of the compound.


The present invention also provides the use of a compound of formula (I) or (II), or a pharmaceutically acceptable salt thereof in the preparation of a medicament for use in the treatment of emesis, anxiety and/or IBS in mammals, including humans. Such treatment may be carried out in the manner or hereinbefore described.

The present invention further provides a pharmaceutical composition for use in the treatment of emesis, anxiety and/or IBS which comprises an effective amount of a compound of formula (I) or (II), or a pharmaceutically acceptable salt thereof and a pharmaceutically acceptable carrier. Such compositions may be prepared in the manner as hereinbefore described.

The following examples illustrate the preparation of compounds of formula (I) and (II) and the following pharmacological data illustrate the invention.

Example 1

3,5-Dichloro-[3 α -(8-methyl-8-azabicyclo(3.2.1)octyl)]benzoate



3,5-Dichlorobenzoic acid (4.78g) was heated at reflux with thionyl chloride (40ml) in dry benzene (100ml) for 2hrs. The mixture was evaporated in vacuo. The residue was treated with dry benzene (50ml) and the mixture was re-evaporated in vacuo. This procedure was carried out three times. An infra-red spectrum of the residue showed complete conversion of the acid to the acid chloride.

Tropine (3.53g) in dry tetrahydrofuran (50ml) was cooled in an ice bath and treated dropwise with butyl lithium (15.7ml; 1.6M solution in hexane), under nitrogen. The mixture was left at ice-bath temperature for 30 minutes then it was evaporated in vacuo (<30°C). The residue was treated with dry tetrahydrofuran (50ml) and cooled in an ice-bath.

The acid chloride (prepared above) in dry tetrahydrofuran (30ml) was added dropwise and the mixture was stirred at room temperature overnight, under nitrogen. The mixture was diluted with dichloromethane and the organic phase was washed with saturated sodium carbonate solution and with brine, dried (Na₂SO₄) and evaporated to give a yellow gum (6g). This was chromatographed on silica gel (120g) using chloroform, 2% methanol in chloroform, 5% methanol in chloroform then 5% methanol in chloroform containing 0.5% ammonium hydroxide, as successive eluants. The resulting pale yellow solid (3.1g) was recrystallised from ethyl acetate to give pale yellow crystals (1st crop 1.55g).


A portion (1.05g) was dissolved in dry ethanol (1.5ml) and treated dropwise with ethanolic hydrogen

chloride (3.1ml of 1.1 molar solution). Addition of a few drops of dry ether gave the hydrochloride salt of the title compound as white crystals (970mg) m.p. 259-61°C.

m.s. $C_{15}H_{17}NO_2Cl_2$ required	313.0636
found	313.0645

Example 2

Indole-[3 α -(8-methyl-8-azabicyclo(3.2.1)octyl)]-3-carboxylate




Tropine (6.35g) in dry tetrahydrofuran (20ml) was cooled in an ice-bath and treated dropwise with butyl lithium (21.3ml of 1.6M solution in hexane). The mixture was stirred for 30 minutes at ice-bath temperature then it was evaporated in vacuo (<30°C). The residue was treated with dry tetrahydrofuran (40ml) and cooled in an ice-bath.

Indole-3-carboxylic acid chloride [prepared from 5g of acid as in Example 1] in dry tetrahydrofuran (20ml) was added dropwise and the resulting mixture was stirred at room temperature, under nitrogen, overnight. The mixture was diluted with dichloromethane and the organic phase was washed with saturated potassium carbonate and with brine, dried (Na_2SO_4) and evaporated to give a yellow solid (8.1g). This was chromatographed on silica gel (250g) using dichloromethane/methanol/ammonium hydroxide (90/10/0.25) then (90/10/0.5) as eluants to give a white solid (3.93g). A portion (2.93g) was recrystallised from ethyl acetate to give the title compound (1.57g) as white crystals. m.p. 199-202°C

$C_{17}H_{26}N_2O_2$ required:	C, 71.81;	H, 7.09;	N, 9.85%
found :	C, 71.74;	H, 7.16;	N, 9.75%

Example 3

3,5-Dimethyl-[3 α -(8-methyl-8-azabicyclo(3.2.1)octyl)]benzoate hydrochloride




HCl.

The title compound was prepared in an analogous manner to the compound of Example 1. m.p. 250-252°C.

Example 4

Indole-[5-(2-methyl-2-azabicyclo(2.2.2)octyl)]-3-carboxylate




Indole 3-carboxylic acid (913mg) was stirred overnight with thionyl chloride (10ml), at room temperature. The excess thionyl chloride was evaporated in vacuo (<30°C) and the residue was washed with dry ether (3 x 50ml).

Meanwhile, 2-methyl-2-azabicyclo(2.2.2)octan-5-ol (800mg) in dry tetrahydrofuran (5ml) at 0°C was treated dropwise with butyl lithium (3.5ml of 1.6M solution in hexane), under nitrogen. The mixture was stirred 30 minutes at ice-bath temperature then the hexane was evaporated in vacuo. The residue was suspended in dry tetrahydrofuran (20ml) and the acid chloride (prepared above) in dry tetrahydrofuran (5ml) was added dropwise. The resulting mixture was stirred at room temperature, under nitrogen, overnight.

The mixture was diluted with dichloromethane and the organic phase was washed with saturated potassium carbonate solution and with brine, dried (Na₂SO₄) and evaporated to give a yellow gum (970mg). This was chromatographed on silica gel (20g) using dichloromethane, 2% methanol in dichloromethane, 5% methanol in dichloromethane then 5% methanol in dichloromethane containing 0.5% ammonium hydroxide as successive eluants to give the title compound (250mg) as a white solid.

Example 5

1,2,3,9-tetrahydro-9-methyl-3-[(2-methyl-1H-imidazol-1-yl)methyl]-4H-carbazol-4-one



The title compound was prepared in accordance with the procedure described in Example 7 of GB 2153821 A m.p. 229-230°C.

Pharmacological Data

Inhibition of Chemotherapy Induced Nausea and Vomiting

Adult male ferrets (initial body weight 1.3-1.8kg) were individually housed, fed once daily (200g Chum Puppy Food plus 50g Lab Diet A) and were supplied with water ad libitum. Additionally, each animal was given $\frac{1}{2}$ pint of milk daily during an acclimatization period.

To facilitate intravenous administration of drugs, a chronic indwelling catheter was surgically implanted into the jugular vein using a modification of the technique described by Florczyk and Schurig, 1981 (Pharmacol. Biochem. Behav., 14, 255-257). Prior to surgery each animal was sedated with ketamine hydrochloride (40mg/animal intramuscularly) and anaesthetised with a halothane-N₂O-O₂ mixture. A four day recovery period was allowed before commencement of an experiment.

For each group of animals, a preliminary study was carried out to establish an intravenous dose level of cytotoxic agent(s) which would give a consistent and reproducible emetic response. The appropriate dose of example was administered intravenously twice to each animal; 30 minutes before and 30-45 minutes after dosing with cytotoxic agent(s). Running controls received vehicle and cytotoxic agent(s) only.

A single emetic response commenced when an animal assumed a characteristic posture with retching and was concluded when either vomitus was expelled or was present in the mouth as evidenced by a chewing movement. The total number of emetic responses was determined during the four hour period following administration of cytotoxic agent(s) and the number of animals completely protected from emesis determined for each treatment.

All solutions were prepared in water for injection. B.P. Cisplatin was prepared from vials of Neoplatin * for injection (Mead Johnson; dose volume of 3ml/kg). Doxorubicin (Adriamycin *); Farmitalia Carlo Erba Ltd. Cyclophosphamide (Endoxana *); W.B. Pharmaceuticals Ltd.

The results obtained are shown below:

Treatment Dose mg/kg i.v. Administered twice	Dose of Cisplatin mg/kg i.v.	No. of animals completely protected/No. of animals used
Vehicle	10	1/11
Compound 1		
0.5	10	3/4
Compound 2		
0.005	10	2/4
0.05		
0.5	10	3/4
Compound 3		
0.5	10	2/3


Compounds 4 and 5

At a treatment dose of 0.5mg/kg of test compound, 6mg/kg of doxorubicin and 80 mg/kg of cyclophosphamide, the number of animals completely protected was 2/3 and 2/2 for compounds 4 and 5 respectively.

Other compounds within formulae (I) and (II) as described in UK Patent Applications 2100259, 2152049, 2125398 and 2153821 may be tested and found to be active in the above test.

Claims

- The use of a compound of formula (II) or a pharmaceutically acceptable salt thereof:




wherein

R_{10} is hydrogen, C_{1-10} alkyl, C_{3-7} cycloalkyl, C_{3-6} alkenyl, phenyl or phenyl- C_{1-3} alkyl; and one of the groups represented by R_{11} , R_{12} and

R_{13} is hydrogen, C_{1-6} alkyl, C_{3-7} cycloalkyl, C_{2-6} alkenyl or phenyl- C_{1-3} alkyl and each of the other groups, which may be the same or different, is hydrogen or C_{1-6} alkyl;

in the manufacture of a medicament for use in the treatment of cytotoxic agent or radiation induced nausea and vomiting, or motion sickness.

- A use according to claim 1 wherein the compound of formula (II) is of formula (IIA):



(IIA)

10 wherein

R_{10}^1 represents a hydrogen atom or a methyl, ethyl, propyl, prop-2-yl, prop-2-enyl or cyclopentyl group;

R_{12}^1 represents a hydrogen atom; and

either R_{11}^1 represents a methyl, ethyl, propyl or prop-2-yl group and R_{13}^1 represents a hydrogen atom; or

R_{11}^1 represents a hydrogen atom and R_{13}^1 represents a methyl or ethyl group.

15 3. A use according to claim 2 wherein R_{10}^1 is methyl and R_{12}^1 and R_{13}^1 are both hydrogen.

4. A use according to claim 1 wherein the compound of formula (II) is 1,2,3,9-tetrahydro-9-methyl-3-[(2-methyl-1H-imidazol-1-yl)methyl]-4H-carbazol-4-one or a pharmaceutically acceptable salt thereof.

5. A use according to any one of claims 1 to 4 wherein the medicament is for use in the treatment of cytotoxic agent induced nausea and vomiting.

20 6. A use according to claim 5 wherein cytotoxic agent is cisplatin, doxorubicin or cyclophosphamide.

7. A use according to claim 6 wherein the cytotoxic agent is doxorubicin or cyclophosphamide.

8. A use according to claim 6 wherein the cytotoxic agent is cisplatin, the compound of formula (II) being administered before or after cisplatin administration.

9. A use according to claim 6 wherein the cytotoxic agent is cisplatin administered intravenously.

25 10. A use according to any one of claims 1 to 4, wherein the medicament is for use in the treatment of radiation induced nausea and vomiting.

11. A use according to any one of claims 1 to 4, wherein the medicament is for use in the treatment of motion sickness.

30

35

40

45

50

55